

Table XV-16
LUNG AND BRONCHUS (Invasive)

SEER INCIDENCE AND U.S. MORTALITY
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity and Sex

SEER Incidence

RACE/ETHNICITY	Rate 2000-2003 ^b Rate per 100,000 persons			Trend 1994-2003 ^c APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	64.8	82.1	52.3	-1.2*	-2.0*	-0.3
White	66.0	81.7	54.7	-1.1*	-2.0*	-0.2
White Hispanic ^d	33.3	45.5	24.5	-1.8*	-2.2*	-1.4*
White Non-Hispanic ^d	67.3	81.9	56.8	-1.0*	-2.0*	-0.1
Black	76.9	112.2	53.1	-1.3*	-2.4*	0.3
Asian/Pacific Islander	39.6	55.7	27.3	-0.8*	-1.0*	-0.2
Amer Ind/Alaska Nat ^e	43.1	55.5	33.8	-3.9*	-5.2*	-1.6
Hispanic ^d	32.6	44.7	24.0	-1.9*	-2.2*	-1.5*

U.S. Mortality^f

RACE/ETHNICITY	Rate 2000-2003 Rate per 100,000 persons			Trend 1994-2003 APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	55.1	74.2	41.2	-0.9*	-1.9*	0.4*
White	55.3	73.4	42.2	-0.7*	-1.8*	0.5*
White Hispanic ^d	24.7	37.8	15.3	-1.3*	-2.3*	0.1
White Non-Hispanic ^d	57.7	76.2	44.3	-0.6*	-1.7*	0.6*
Black	62.5	97.2	39.8	-1.6*	-2.7*	0.3
Asian/Pacific Islander	27.2	38.6	18.6	-1.3*	-1.6*	-0.7
Amer Ind/Alaska Nat	33.1	41.4	26.8	-2.0*	-3.8*	0.3
Hispanic ^d	23.9	36.6	14.7	-1.3*	-2.2*	0.1

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
- ^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- ^d Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. The 2000-2003 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1994-2003 Hispanic and Non-Hispanic mortality trends exclude deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.
- ^e Incidence data for American Indians/Alaska Natives include cases from Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, Atlanta, and the Alaska Native Registry. Incidence rates are for diagnosis years 1999-2002. Incidence trends are for diagnosis years 1994-2002.
- ^f Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- * The APC is significantly different from zero (p<.05).